

SUPPLEMENTARY TABLE. Age-adjusted prevalence of binge drinking among adults aged ≥18 years, by state — Behavioral Risk Factor Surveillance System, United States,* 2018

State	Weighted number of adults reporting binge drinking[†] (x1000)	Binge drinking[†] prevalence[§], % (95% CI)
Alabama	445	13.6 (12.3–15.0)
Alaska	83	16.0 (14.0–18.3)
Arizona	791	16.7 (15.2–18.3)
Arkansas	313	16.0 (14.3–17.9)
California	4,486	16.6 (15.7–17.5)
Colorado	771	19.8 (18.7–20.9)
Connecticut	445	18.7 (17.4–20.0)
Delaware	117	18.3 (16.7–20.1)
District of Columbia	128	22.5 (20.7–24.3)
Florida	2,501	17.5 (16.2–19.0)
Georgia	1,065	15.1 (14.1–16.1)
Hawaii	201	20.9 (19.5–22.3)
Idaho	186	16.0 (14.2–17.8)
Illinois	1,827	20.5 (19.1–22.0)
Indiana	772	17.1 (15.9–18.5)
Iowa	523	24.5 (23.4–25.6)
Kansas	324	16.7 (15.7–17.8)
Kentucky	486	15.8 (14.4–17.4)
Louisiana	560	18.1 (16.5–19.7)
Maine	173	19.5 (17.9–21.2)
Maryland	594	14.5 (13.6–15.5)
Massachusetts	1,003	21.1 (19.6–22.6)
Michigan	1,340	19.8 (18.7–20.9)
Minnesota	835	21.7 (20.9–22.5)
Mississippi	273	13.7 (12.3–15.2)
Missouri	794	18.9 (17.3–20.6)
Montana	141	19.7 (18.1–21.5)
Nebraska	292	22.5 (21.4–23.7)
Nevada	333	16.0 (14.1–18.1)
New Hampshire	166	18.3 (16.6–20.2)
New Jersey	866	14.4 (12.4–16.6)
New Mexico	220	15.6 (14.2–17.1)
New York	2,271	17.6 (16.8–18.4)
North Carolina	1,084	15.8 (14.4–17.4)
North Dakota	121	22.7 (21.1–24.5)
Ohio	1,372	17.2 (16.0–18.4)
Oklahoma	362	13.6 (12.3–15.1)
Oregon	482	17.0 (15.8–18.3)
Pennsylvania	1,610	18.5 (17.2–19.9)
Rhode Island	129	17.8 (16.1–19.6)
South Carolina	569	16.3 (15.1–17.5)
South Dakota	133	22.3 (20.3–24.4)
Tennessee	700	15.7 (14.1–17.4)
Texas	3,358	17.6 (16.1–19.2)
Utah	228	10.5 (9.8–11.3)
Vermont	79	18.5 (16.9–20.2)
Virginia	967	16.4 (15.3–17.6)
Washington	779	15.4 (14.5–16.4)
West Virginia	167	13.9 (12.5–15.4)

Wisconsin	1,008	25.8 (24.1–27.7)
Wyoming	74	18.5 (16.9–20.2)

Abbreviation: CI=confidence interval.

* The study sample included 398,485 respondents from all 50 states and the District of Columbia with complete information on age, sex, and binge drinking. Weighted numbers were derived through application of survey weights.

† Binge drinking was defined as consuming ≥ 5 drinks for men or ≥ 4 drinks for women on an occasion during the past 30 days.

§ Prevalence estimates for states were age-standardized to the 2000 U.S. Census population.